

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 August 2004 (26.08.2004)

PCT

(10) International Publication Number
WO 2004/071428 A2

(51) International Patent Classification⁷: **A61K**
(21) International Application Number:
PCT/US2004/003433

(22) International Filing Date: 6 February 2004 (06.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/365,369 11 February 2003 (11.02.2003) US
10/772,720 4 February 2004 (04.02.2004) US

(71) Applicant (for all designated States except US): **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DONDE, Yariv** [US/US]; 24386 Antilles Way, Dana Point, CA 92629 (US). **NGUYEN, Jeremiah, H.** [US/US]; 16146 Bamboo Street, La Puente, CA 91744 (US).

(74) Agents: **JOHNSON, Brent, A.** et al.; Allergan, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— without international search report and to be republished upon receipt of that report

[Continued on next page]


(54) Title: 10,10-DIALKYL PROSTANOIC ACID DERIVATIVES AS AGENTS FOR LOWERING INTRAOCULAR PRESSURE

(SCHEME 1)


EQUATION 1



EQUATION 2



EQUATION 3



(57) Abstract: The present invention provides a method of treating ocular hypertension or glaucoma which comprises administering to an animal having ocular hypertension or glaucoma therapeutically effective amount of a compound represented by the general formula (I); wherein the dashed line indicates the presence or absence of a bond, the hatched wedge indicates the α (down) configuration, and the solid triangle indicates the β (up) configuration; B is a single, double, or triple covalent bond; n is 0-6; X is CH₂, S or O; Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂, CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, CH₂OR, P(O)(OR)₂, CONRSO₂R, SONR₂, or formula (Ia) R is H, C₁₋₆ alkyl or C₂₋₆ alkenyl; R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

10,10-DIALKYL PROSTANOIC ACID DERIVATIVES AS AGENTS FOR LOWERING INTRAOCULAR PRESSURE

Field of the Invention

5

The present invention relates to prostanoid acid derivatives as potent ocular hypotensives that are particularly suited for the management of glaucoma.

Background of the Invention

10

Description of Related Art

Ocular hypotensive agents are useful in the treatment of a number of various ocular hypertensive conditions, such as post-surgical and post-laser trabeculectomy ocular hypertensive episodes, glaucoma, and as presurgical adjuncts.


Glaucoma is a disease of the eye characterized by increased intraocular pressure. On the basis of its etiology, glaucoma has been classified as primary or secondary. For example, primary glaucoma in adults (congenital glaucoma) may be either open-angle or acute or chronic angle-closure. Secondary glaucoma results from pre-existing ocular diseases such as uveitis, intraocular tumor or an enlarged cataract.

The underlying causes of primary glaucoma are not yet known. The increased intraocular tension is due to the obstruction of aqueous humor outflow. In chronic open-angle glaucoma, the anterior chamber and its anatomic structures appear normal, but drainage of the aqueous humor is impeded. In acute or chronic angle-closure glaucoma, the anterior chamber is shallow, the filtration angle is narrowed, and the iris may obstruct the trabecular meshwork at the entrance of the canal of Schlemm. Dilation of the pupil may push the root of the iris forward against the angle, and may produce pupillary block and thus precipitate an acute attack. Eyes with narrow anterior chamber angles are predisposed to acute angle-closure glaucoma attacks of various degrees of severity.

Secondary glaucoma is caused by any interference with the flow of aqueous humor from the posterior chamber into the anterior chamber and subsequently, into the canal of Schlemm. Inflammatory disease of the anterior segment may prevent aqueous escape by causing complete posterior synechia in
5 iris bombe, and may plug the drainage channel with exudates. Other common causes are intraocular tumors, enlarged cataracts, central retinal vein occlusion, trauma to the eye, operative procedures and intraocular hemorrhage.

Considering all types together, glaucoma occurs in about 2% of all persons over the age of 40 and may be asymptotic for years before progressing to
10 rapid loss of vision. In cases where surgery is not indicated, topical β -adrenoreceptor antagonists have traditionally been the drugs of choice for treating glaucoma.

Certain eicosanoids and their derivatives have been reported to possess ocular hypotensive activity, and have been recommended for use in glaucoma
15 management. Eicosanoids and derivatives include numerous biologically important compounds such as prostaglandins and their derivatives. Prostaglandins can be described as derivatives of prostanoic acid which have the following structural formula:



Various types of prostaglandins are known, depending on the structure and substituents carried on the alicyclic ring of the prostanoic acid skeleton. Further classification is based on the number of unsaturated bonds in the side chain indicated by numerical subscripts after the generic type of prostaglandin
25 [e.g. prostaglandin E₁ (PGE₁), prostaglandin E₂ (PGE₂)], and on the

configuration of the substituents on the alicyclic ring indicated by α or β [e.g. prostaglandin $F_2\alpha$ ($PGF_2\beta$)].

Prostaglandins were earlier regarded as potent ocular hypertensives, however, evidence accumulated in the last decade shows that some
5 prostaglandins are highly effective ocular hypotensive agents, and are ideally suited for the long-term medical management of glaucoma (see, for example, Bito, L.Z. Biological Protection with Prostaglandins, Cohen, M.M., ed., Boca Raton, Fla, CRC Press Inc., 1985, pp. 231-252; and Bito, L.Z., Applied Pharmacology in the Medical Treatment of Glaucomas Drance, S.M. and
10 Neufeld, A.H. eds., New York, Grune & Stratton, 1984, pp. 477-505. Such prostaglandins include $PGF_2\alpha$, $PGF_1\alpha$, PGE_2 , and certain lipid-soluble esters, such as C_1 to C_2 alkyl esters, e.g. 1-isopropyl ester, of such compounds.

Although the precise mechanism is not yet known experimental results indicate that the prostaglandin-induced reduction in intraocular pressure results
15 from increased uveoscleral outflow [Nilsson et. al., Invest. Ophthalmol. Vis. Sci. (suppl), 284 (1987)].

The isopropyl ester of $PGF_2\alpha$ has been shown to have significantly greater hypotensive potency than the parent compound, presumably as a result of its more effective penetration through the cornea. In 1987, this compound was
20 described as "the most potent ocular hypotensive agent ever reported" [see, for example, Bito, L.Z., Arch. Ophthalmol. 105, 1036 (1987), and Siebold et al., Prodrug 5 3 (1989)].

Whereas prostaglandins appear to be devoid of significant intraocular side effects, ocular surface (conjunctival) hyperemia and foreign-body sensation have
25 been consistently associated with the topical ocular use of such compounds, in particular $PGF_2\alpha$ and its prodrugs, e.g., its 1-isopropyl ester, in humans. The clinical potentials of prostaglandins in the management of conditions associated with increased ocular pressure, e.g. glaucoma are greatly limited by these side effects.

30 In a series of co-pending United States patent applications assigned to Allergan, Inc. prostaglandin esters with increased ocular hypotensive activity

accompanied with no or substantially reduced side-effects are disclosed. The co-pending USSN 596,430 (filed 10 October 1990, now U.S. Patent 5,446,041), relates to certain 11-acyl-prostaglandins, such as 11-pivaloyl, 11-acetyl, 11-isobutyryl, 11-valeryl, and 11-isovaleryl PGF₂α. Intraocular pressure reducing
5 15-acyl prostaglandins are disclosed in the co-pending application USSN 175,476 (filed 29 December 1993). Similarly, 11,15- 9,15 and 9,11-diesters of prostaglandins, for example 11,15-dipivaloyl PGF₂α are known to have ocular hypotensive activity. See the co-pending patent applications USSN Nos. 385,645 (filed 07 July 1989, now U.S. Patent 4,994,274), 584,370 (filed 18 September
10 1990, now U.S. Patent 5,028,624) and 585,284 (filed 18 September 1990, now U.S. Patent 5,034,413). Recently, we have also shown that 17-naphthyl and benzothienyl prostaglandin compounds also have ocular hypotensive activity (USSN 859,770, filed 17 May 2001). The disclosures of all of these patent applications are hereby expressly incorporated by reference.


15 Certain 15,15-dimethyl prostaglandins with antihypertensive, gastric acid secretion inhibition, and smooth muscle stimulant properties, are known to have improved metabolic stability. These are described by Pernet *et al* in US Patent 4,117,014 (filed 23 December 1976), the disclosure of which is hereby expressly incorporated by reference.

20

Summary of the Invention

The present invention concerns a method of treating ocular hypertension which comprises administering to a mammal having ocular hypertension a
25 therapeutically effective amount of a compound of Formula I

5

**Formula I**


wherein the dashed line indicates the presence or absence of a bond, the hatched
 5 wedge indicates the α (down) configuration, and the solid triangle indicates the
 β (up) configuration;

B is a single, double, or triple covalent bond;

n is 0-6;

X is CH₂, S or O;

- 10 Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂,
 CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, CH₂OR, P(O)(OR)₂, CONRSO₂R,
 SONR₂, or



R is H, C₁₋₆ alkyl or C₂₋₆ alkenyl;

- 15 R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be
 bonded to each other such that they form a ring incorporating the carbon to
 which they are commonly attached;

R⁴ is hydrogen, R, C(=O)R, or any group that is easily removed under
 physiological conditions such that R⁴ is effectively hydrogen;

- 20 R⁵ is hydrogen or R;

R⁶ is

- i) hydrogen;

- ii) a linear or branched hydrocarbon containing between 1 and 8 carbon atoms, which may contain one or more double or triple bonds, or oxygen or halogen derivatives of said hydrocarbon, wherein 1-3 carbon or hydrogen atoms may be substituted by O or a halogen; or
- 5 iii) aryloxy, heteroaryloxy, C₃₋₈ cycloalkyloxy, C₃₋₈ cycloalkyl, C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₆₋₁₀ aryl, C₃₋₁₀ heteroaryl, aryloxy, heteroaryloxy, C₁₋₆ alkyl,
- 10 OR, SR, and SO₂R.

In another aspect, the present invention relates to a pharmaceutical product, comprising a container adapted to dispense its contents in a metered form; and an ophthalmic solution therein, as hereinabove defined.

15 In another aspect, certain of the compounds represented by the above formula, disclosed below and utilized in the method of the present invention are novel and unobvious.

In a further aspect, certain elements of the processes of preparing the compounds represented by the above formula and described herein are novel and unobvious.

20

Brief Description of the Drawing Figures

Schemes 1-8 illustrate possible ways to prepare compounds of the invention.


25

Detailed Description of the Invention


The present invention relates to the use of prostanoic acid derivatives as ocular hypotensives. The compounds used in accordance with the present

30 invention are encompassed by the following structural formula I:

7

**Formula I**

A preferred group of the compounds of the present invention includes compounds that do not have the following structural formula II:

**Formula II**

5

wherein A is CO₂H, CO₂Me, or CO₂Et;

D is a single, double, or triple covalent bond;

E is a linear, branched, or cycloalkyl chain of 3 to 7 carbons,

trifluoromethylbutyl, hydroxylalkyl, or CH₂R⁷ wherein R⁷ is phenyl,

10 cyclopentyl, phenoxy, chlorophenoxy, propoxy, or -CH₂SCH₂CH₃;

J is hydrogen, R, C(=O)R, or any group that is easily removed under physiological conditions such that R⁴ is effectively hydrogen; and

G is H or CH₃.


As used herein, the symbols "Me" and "Et" refer to the moieties

15 commonly referred to as "methyl" and "ethyl" by those of ordinary skill in the art.

In other compounds related to Formula I and Formula II, A is CO_2R^8 , wherein R^8 is any linear, branched, or cyclic alkyl group having from 3 to 6 carbons.


Another preferred group includes compounds having formula III:

5




Formula III

Another preferred group includes compounds having formula IV:



Formula IV

Another preferred group includes compounds having formula V:



Formula V

wherein at least one of R^2 and R^3 is not methyl.

In the above formulae, the substituents and symbols are as hereinabove defined.


In the above formulae:

- 5 Preferably Y is any pharmaceutically acceptable salt of CO_2H or CO_2R .
More preferably Y is CO_2H or CO_2Me .

Preferably n is 2.

- Preferably, R^6 is C_{6-10} aryl or C_{3-10} heteroaryl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R . More preferably R^6 is phenyl, naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R . Most preferred is 3-chlorobenzothien-2-yl.

- 15 Another preferred group includes compounds having formula XIII:





Formula XIII

wherein B represents a single or double bond;

- and R^6 is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R .

- In another aspect of this invention, certain elements of the method making the compounds of the invention are novel and unobvious. One such novel and unobvious element is the application of the use of Baker's yeast as a reducing agent as reported by Brooks and coworkers (Brooks, *et. al.*,
25 "Asymmetric Microbial Reduction of Prochiral 2,2-Disubstituted

Cycloalkanediones", *J. Org. Chem.*, **1987**, 52, 3223-3232) in the synthesis of compounds of this invention. In this novel and unobvious application of this reaction, Baker's yeast is used to carry out an asymmetric reduction of a compound of formula VII, which is a 2,2-dialkylcyclopentane-1,3-dione, to a
5 compound of formula VIII, which is a 2,2-dialkyl-3(S)-hydroxycyclopentanone. A compound of formula VIII is then used to prepare compounds of this invention.


**Formula VI****Formula VII**

The two alkyl groups, R^2 and R^3 of the compounds of formula VI and VII, in
10 this reaction are the same as those defined for compounds of Formula I above. In the case where the two alkyl groups are different, a mixture of diastereomers is formed, which can be separated by conventional separation methods to obtain the enantiomerically pure products.

Preparation of 2,2-dialkylcyclopentan-1,3-diones is well known in the
15 art. One convenient way that a large variety of these compounds can be prepared is by base-mediated alkylation of carbon-2 of the cyclopentane-1,3-dione using an alkyl halide or equivalent compound. This type of reaction is well known in the art. The preparation of three general types of 2,2-dialkylcyclopentan-1,3-diones using this alkylation reaction is illustrated in
20 Scheme 1. Compounds where one of the alkyl groups is methyl can be prepared by a simple alkylation reaction from commercially available 2-methylcyclopentan-1,3-dione **1** (Equation 1). In the case neither of the alkyl groups in the 2,2-dialkylcyclopentan-1,3-dione are methyl (compound **2b**), these compounds can be prepared from cyclopentan-1,3-dione by two consecutive
25 alkylation reactions (Equation 2). In the case where the two alkyl groups in the 2,2-dialkylcyclopentan-1,3-dione are the same, these alkylation reactions can be


carried out in a one-pot procedure. In the case where the two alkyl groups to form a cyclic compound incorporating C₂ of the cyclopentanone into the ring, otherwise known as a spiroketone, these compounds can be prepared by using a dihaloalkane or equivalent compound to carry out an intermolecular alkylation followed by an intramolecular alkylation (Equation 3), which could be carried out in a one or two pot process. Those skilled in the art will recognize that there are many ways to prepare 2,2-dialkylcyclopentan-1,3-diones, and the reactions of Scheme 1 are included to illustrate that these compounds can be readily prepared or obtained by those skilled in the art, and are not intended to limit the scope of the invention in any way.

In another novel and unobvious aspect of this invention, compounds of the invention represented by Formula VIII are prepared by a process that comprises the following steps:




Formula VIII

15 i) reacting a compound of Formula IX with a compound of Formula X in the presence of a suitable base to form a compound of Formula XI;




Formula IX




Formula X

12

**Formula XI**

ii) coupling a compound of Formula XI with a compound of Formula XII; and

**Formula XII**

5 iii) removing the protecting groups and separating the diastereomers to obtain the desired products;

wherein the hatched wedges indicate the α (down) configuration, the solid triangles indicate the β (up) configuration, and the wavy lines indicate either the cis (Z) or trans (Z) conformation;

10 n is 0-6;

B is a single, double, or triple covalent bond;

J is a protecting group that can be easily removed to form the respective hydroxide group without affecting the rest of the molecule;

R is C₁₋₆ alkyl or C₂₋₆ alkenyl;

15 R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached;

X is S or O; and

M is a group that comprises one or more metal atoms.















20 All of the compounds encompassed by this invention can be prepared using the methods described above supplemented by methods known to those skilled in the art. The synthesis of several compounds of the invention is illustrated in

Schemes 2-7. These Schemes that are included herein are merely illustrative and are not intended to limit the scope of the invention in any way. Although there are several ways the reduction of compounds of Formula VI to Formula VII could be incorporated into the synthesis of these compounds, one

5 convenient way to this is shown in Scheme 2. In this Scheme, compound **2** is a compound of Formula VI and compound **3** is a compound of formula VII. However, those skilled in the art will recognize that there are many ways in which the reduction could be used to prepare compounds of this invention.

Table 1

Structure	Low Rf diastereomer	High Rf diastereomer
	21	22
	23	24
	34	35
	36	37
	38	39
	40	41
		42
		43
		44
		45
	46	47

	48	49
	50	51
	52	53
	54	55
	56	57
	58	59
	60	61
	62	63
	64	65
	66	67
	68	69
	70	71
	72	73
	74	75

The compounds named below, and illustrated in Table 1, are especially preferred representatives of the compounds of the present invention:

- (3-{(1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl}-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid methyl ester (**21, 22**);
- (3-{(1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl}-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid
- 5 (23, 24);
- (*Z*)-7-{(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl}-hept-5-ynoic acid methyl ester (**34, 35**);
- 10 (*Z*)-7-{(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl}-hept-5-ynoic acid (**36,37**);
- (*Z*)-7-{(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl}-hept-5-enoic acid methyl ester (**38,39**);
- 15 (*Z*)-7-{(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl}-hept-5-enoic acid (**40,41**);
- 7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**42**)
- 7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-
- 20 oxo-cyclopentyl]-hept-5-ynoic acid (**43**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**44**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**45**)
- 25 (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-3-hydroxy-4-phenyl-but-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**46, 47**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-3-hydroxy-4-phenyl-but-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**48, 49**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-
- 30 dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**50,51**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-4-Hydroxy-5-((*E*)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**52,53**)

- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**54,55**)
- 7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-heptanoic acid (**56,57**)
- 5 (*Z*)-7-[(1*R*,4*S*,5*R*)-5-(4-Benzo[*b*]thiophen-2-yl-3-hydroxy-butyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**58,59**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid ethylamide (**60,61**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-
- 10 hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid diethylamide (**62,63**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (2-hydroxy-ethyl)-amide (**64,65**)
- 15 (3*S*,4*R*,5*R*)-4-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-3-hydroxy-2,2-dimethyl-5-[(*Z*)-6-(1-*H*-tetrazol-5-yl)-hex-2-enyl]-cyclopentanone (**66,67**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid amide (**68,69**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-
- 20 hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**70,71**)
- 7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**72,73**)
- 7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**74,75**)
- 25 Pharmaceutical compositions may be prepared by combining a therapeutically effective amount of at least one compound according to the present invention, or a pharmaceutically acceptable acid addition salt thereof, as an active ingredient, with conventional ophthalmically acceptable pharmaceutical excipients, and by preparation of unit dosage forms suitable for topical ocular use.
- 30 The therapeutically efficient amount typically is between about 0.0001 and about 5% (w/v), preferably about 0.001 to about 1.0% (w/v) in liquid formulations.

For ophthalmic application, preferably solutions are prepared using a physiological saline solution as a major vehicle. The pH of such ophthalmic solutions should preferably be maintained between 6.5 and 7.2 with an appropriate buffer system. The formulations may also contain conventional,
5 pharmaceutically acceptable preservatives, stabilizers and surfactants.

Preferred preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A preferred surfactant is, for example, Tween 80.
10 Likewise, various preferred vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

Tonicity adjustors may be added as needed or convenient. They include,
15 but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include
20 acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated
25 hydroxytoluene.

Other excipient components which may be included in the ophthalmic preparations are chelating agents. The preferred chelating agent is edentate disodium, although other chelating agents may also be used in place or in conjunction with it.

30 The ingredients are usually used in the following amounts:

<u>Ingredient</u>	<u>Amount (% w/v)</u>
active ingredient	about 0.001-5

18

	preservative	0-0.10
	vehicle	0-40
	tonicity adjustor	1-10
	buffer	0.01-10
5	pH adjustor	q.s. pH 4.5-7.5
	antioxidant	as needed
	surfactant	as needed
	purified water	as needed to make 100%

10 The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

 The ophthalmic formulations of the present invention are conveniently packaged in forms suitable for metered application, such as in containers
15 equipped with a dropper, to facilitate the application to the eye. Containers suitable for dropwise application are usually made of suitable inert, non-toxic plastic material, and generally contain between about 0.5 and about 15 ml solution.

Synthetic Examples

20 The methods of preparing compounds of this invention are further illustrated by the following non-limiting Examples, which are summarized in the reaction schemes of Figures 1-7 wherein the compounds are identified by the same designator in both the Examples and the Figures.

2-Alkyl-cyclopentane-1,3-dione (1a). A mixture of 1,3-
25 cyclopentanedione (89.4 mmol, Aldrich), I-R² (96.4 mmol, Aldrich), and KOH (5.097 g, 90.8 mmol) in H₂O (25 mL)/ dioxane (75 mL) is heated at reflux. After 5 h, a solution of KOH (2 g) and I-R² (2 mmol) in H₂O (5 mL)/ dioxane (15 mL) is added and after another 3 h at reflux the solution is allowed to stir at room temperature overnight. In the morning, the reaction is continued by
30 addition of a solution of KOH (2 g) and I-R² (2.4 mmol) in H₂O (5 mL)/ dioxane (15 mL) and heating at reflux. After 4 h, the mixture is allowed to cool

to room temperature and is extracted with ether (1 x 100mL, 3 x 75 mL). The combined ether extracts are evaporated, the residue is combined with HCl (50 mL 10%), and the resulting mixture is placed in a 120 °C oil bath until boiling is observed (ca. 15 min.). The mixture is then allowed to cool to room

- 5 temperature, is neutralized by addition of NaHCO₃ solution (150 mL, saturated) and the resulting mixture is then extracted with CH₂Cl₂ (4 x 75 mL). The combined CH₂Cl₂ solution is dried (MgSO₄), filtered and evaporated to leave a brown oil which is used directly in the next step.

- 2-Alkyl-2-methyl-cyclopentane-1,3-dione (2a).** A mixture of 2-
10 methyl-1,3-cyclopentanedione (10.025 g, 89.4 mmol, Aldrich), I-R² (96.4 mmol, Aldrich), and KOH (5.097 g, 90.8 mmol) in H₂O (25 mL)/ dioxane (75 mL) is heated at reflux. After 5 h, a solution of KOH (2 g) and I-R² (2 mmol) in H₂O (5 mL)/ dioxane (15 mL) is added and after another 3 h at reflux the solution is allowed to stir at room temperature overnight. In the morning, the reaction is
15 continued by addition of a solution of KOH (2 g) and I-R² (2.4 mmol) in H₂O (5 mL)/ dioxane (15 mL) and heating at reflux. After 4 h, the mixture is allowed to cool to room temperature and is extracted with ether (1 x 100mL, 3 x 75 mL). The combined ether extracts are evaporated, the residue is combined with HCl (50 mL 10%), and the resulting mixture is placed in a 120 °C oil bath until
20 boiling is observed (ca. 15 min.). The mixture is then allowed to cool to room temperature, is neutralized by addition of NaHCO₃ solution (150 mL, saturated) and the resulting mixture is then extracted with CH₂Cl₂ (4 x 75 mL). The combined CH₂Cl₂ solution is dried (MgSO₄), filtered and evaporated to leave a brown oil which is used directly in the next step.

2,2-Dialkyl-methyl-cyclopentane-1,3-dione (2b). A mixture of 2-alkyl-1,3-cyclopentanedione **1a** (89.4 mmol, Aldrich), I-R³ (96.4 mmol, Aldrich), and KOH (5.097 g, 90.8 mmol) in H₂O (25 mL)/ dioxane (75 mL) is heated at reflux. After 5 h, a solution of KOH (2 g) and I-R³ (2 mmol) in H₂O (5 mL)/ dioxane (15 mL) is added and after another 3 h at reflux the solution is allowed to stir at room temperature overnight. In the morning, the reaction is continued by addition of a solution of KOH (2 g) and I-R³ (2.4 mmol) in H₂O (5 mL)/ dioxane (15 mL) and heating at reflux. After 4 h, the mixture is allowed to cool to room temperature and is extracted with ether (1 x 100 mL, 3 x 75 mL). The combined ether extracts are evaporated, the residue is combined with HCl (50 mL 10%), and the resulting mixture is placed in a 120 °C oil bath until boiling is observed (ca. 15 min.). The mixture is then allowed to cool to room temperature, is neutralized by addition of NaHCO₃ solution (150 mL, saturated) and the resulting mixture is then extracted with CH₂Cl₂ (4 x 75 mL). The combined CH₂Cl₂ solution is dried (MgSO₄), filtered and evaporated to leave a brown oil which is used directly in the next step.

Spiro[2.4]heptane-4,7-dione (2c). A mixture of 2-alkyl-1,3-cyclopentanedione **1a** (89.4 mmol, Aldrich), 1,2-dibromoethane (120 mmol, Aldrich), and KOH (5.097 g, 90.8 mmol) in H₂O (25 mL)/ dioxane (75 mL) is heated at reflux for 24 hours. The mixture is allowed to cool, and the crude product is extracted with ether (1 x 100mL, 3 x 75 mL). The combined ether extracts are evaporated, the residue is combined with HCl (50 mL 10%), and the resulting mixture is placed in a 120 °C oil bath until boiling is observed (ca. 15 min.). The mixture is then allowed to cool to room temperature, is neutralized

by addition of NaHCO₃ solution (150 mL, saturated) and the resulting mixture is then extracted with CH₂Cl₂ (4 x 75 mL). The combined CH₂Cl₂ solution is dried (MgSO₄), filtered and evaporated to leave a brown oil which is used directly in the next step.

5 **2,2-Dimethyl-cyclopentane-1,3-dione (2).** The published procedure was followed. (Agosta, W.C.; Smith, A.B. *J.Org.Chem.* **1970**, 35, 3856) A mixture of 2-methyl-1,3-cyclopentanedione (10.025 g, 89.4 mmol, Aldrich), methyl iodide (6.0 mL, 96.4 mmol, Aldrich), and KOH (5.097 g, 90.8 mmol) in H₂O (25 mL)/ dioxane (75 mL) was heated at reflux. After 5 h, a solution of
10 KOH (2 g) and MeI (2.4 mL) in H₂O (5 mL)/ dioxane (15 mL) was added and after another 3 h at reflux the solution was allowed to stir at room temperature overnight. In the morning, the reaction was continued by addition of a solution of KOH (2 g) and MeI (2.4 mL) in H₂O (5 mL)/ dioxane (15 mL) and heating at reflux. After 4 h, the mixture was allowed to cool to room temperature and was
15 extracted with ether (1 x 100mL, 3 x 75 mL). The combined ether extracts were evaporated, the residue combined with HCl (50 mL 10%), and the resulting mixture was placed in a 120 °C oil bath until boiling was observed (ca. 15 min.). The mixture was then allowed to cool to room temperature, was neutralized by addition of NaHCO₃ solution (150 mL, saturated) and the
20 resulting mixture then extracted with CH₂Cl₂ (4 x 75 mL). The combined CH₂Cl₂ solution was dried (MgSO₄), filtered and evaporated to leave a brown oil (10.474 g, 83 mmol, 93%) which was used directly in the next step.

(S)-3-Hydroxy-2,2-dimethyl-cyclopentanone (3). The published procedure was followed. (Brooks, D.W.; Hormoz, M.; Grothaus, P.G. *J.Org.Chem.* **1987**,

52, 3223) A 35 °C (internal temperature) solution of D-glucose (106.73 g, 592 mmol, Aldrich) in H₂O (690 mL) in a 4 L Erlenmeyer was treated with baker's yeast (71.065 g, Fleischmann's). The mixture was allowed to ferment for 2 h, then 2,2-dimethyl-cyclopentane-1,3-dione (**2**) (7.316 g, 58 mmol) was added.

5

The mixture was stirred for 48 h and then filtered through celite, washing with about 1 L CH₂Cl₂. The filtration was difficult due to the thick consistency of the yeast and it helped to continually add CH₂Cl₂ to the mixture and scrape the top of the celite layer with a spatula. The filtrate was transferred to a separatory funnel, and 100 mL brine was added and the layers were separated. Brine (400 mL) was added to the aqueous layer and the resulting solution extracted further with CH₂Cl₂ (3 x 500 mL). The combined CH₂Cl₂ solution was dried (MgSO₄), filtered and evaporated to leave a yellow oil. Flash chromatography (11 x 5 cm, 20% EtOAc/hexs → 25% → 30% → 40% → 50%) gave alcohol **3** (2.435 g, 19 mmol, 33%).

15

The enantiomeric excess of **3** was assayed by ¹H NMR of the corresponding Mosher's ester which was prepared by treatment of alcohol **3** (11 mg, 0.09 mmol) in dichloroethane (0.3 mL, Aldrich) with pyridine (27 μL, 0.33 mmol, Aldrich) and (*R*)-α-methoxy-α-trifluoromethylphenylacetic acid chloride (58 μL, 0.31 mmol, Fluka). The mixture was stirred overnight and then partitioned between water (10 mL) and ether (10 mL). The ether layer was washed with 1 M HCl (10 mL) and saturated NaHCO₃ solution and then was dried (MgSO₄), filtered and evaporated. ¹H NMR analysis was done on the crude ester.

20

(*S*)-3-(*tert*)-Butyl-dimethyl-silanyloxy-2,2-dimethyl-cyclopentanone (**4**). A solution of alcohol **3** (520 mg, 4.1 mmol) and 2,6-lutidine (0.56 mL, 4.8 mmol, Aldrich) in CH₂Cl₂ (8.0 mL, Aldrich) was treated with TBSOTf (1.0 mL, 4.3 mmol, Aldrich). After 5.5 h, saturated NaHCO₃ solution (20 mL) was added and
5 the mixture extracted with CH₂Cl₂ (20 mL). The CH₂Cl₂ solution was washed with 20 mL each of 1 M HCl, saturated NaHCO₃ solution, and brine and then was dried (MgSO₄), filtered and evaporated. Flash chromatography (5 x 5 cm, 10% Et₂O/pentane) gave TBS ether **4** (698 mg, 2.9 mmol, 70%).

(*S*)-3-(*tert*)-Butyl-dimethyl-silanyloxy-2,2-dimethyl-5-phenylselenanyl-
10 cyclopentanone (**5**). A solution of TBS ether **4** (1.496 g, 6.2 mmol) in THF (2 mL, Aldrich) was added dropwise to a -78 °C solution of LDA (4.9 mL, 7.3 mmol, 1.5 M/cyclohexane, Aldrich) in THF (22 mL, Aldrich), rinsing with 2 mL THF. After 15 min., a solution of PhSeCl (1.424 g, 7.4 mmol, Aldrich) in THF (2 mL) was quickly added by cannula, rinsing with 2 mL THF. The
15 solution was stirred for 10 min. and then partitioned between 50 mL 0.5 M HCl and 75 mL ether. The ether layer was washed with 30 mL each of water, saturated NaHCO₃ solution, and brine and then was dried (MgSO₄), filtered and evaporated. Flash chromatography (2% EtOAc/hexs → 4%) gave phenylselenide **5** (1.641 g, 4.1 mmol, 67%) along with 476 mg of mixed
20 fractions containing a lower R_f impurity.

(*S*)-4-(*tert*)-Butyl-dimethyl-silanyloxy-5,5-dimethyl-cyclopent-2-enone(**6**). A solution of selenide **5** (1.641 g, 4.1 mmol) and pyridine (0.62 mL, 7.7 mmol, Aldrich) in CH₂Cl₂ (13 mL, Aldrich) was treated with H₂O (1 mL) and 30% H₂O₂ (1.1 mL, Aldrich). The mixture was stirred for 30 min. and then was

partitioned between 25 mL CH₂Cl₂ and 25 mL saturated NaHCO₃ solution. The aqueous layer was extracted with 25 mL CH₂Cl₂ and the combined CH₂Cl₂ solution washed with 1 M HCl (2 x 25 mL) and brine (50 mL). The solution was then dried (MgSO₄), filtered and evaporated to leave an orange oil. Flash chromatography (6 x 4 cm, 10% ether/pentane) gave enone **6** (572 mg, 2.4 mmol, 59%).

(3-Mercapto-propylsulfanyl)-acetic acid methyl ester (8). An ice-cold solution of 1,3-dithiane (2.0 mL, 19.9 mmol) in THF (40 mL) was treated with NaH (819 mg, 20.5 mmol). After 30 min., methyl bromoacetate (1.9 mL, 20.0 mmol) was added and the mixture stirred for 3.5 h at room temperature. The reaction was quenched by addition of MeOH and then 50 mL 1 M HCl. The mixture was extracted with ether (2 x 50 mL) and the combined ether solution washed with saturated sodium bicarbonate solution (50 mL) and brine (50 mL) and then was dried (MgSO₄), filtered and evaporated. Purification by flash chromatography on silica gel (10-15% ethyl acetate/hexanes) gave 971 mg (5.38 mmol, 27 %) of the thiol.

{3-[(S)-3-(tert)-Butyl-dimethyl-silanyloxy]-4,4-dimethyl-5-oxo-cyclopent-1-enylsulfanyl]-propylsulfanyl}-acetic acid methyl ester (10). A solution of enone **6** (156 mg, 0.65 mmol) in MeOH (4.3 mL) was treated with 30 % H₂O₂ (0.21 mL) and 1 M NaOH (32 µL). After 4 h, 20 mL saturated ammonium chloride solution was added and the mixture was extracted with dichloromethane (3 x 10 mL). The combined dichloromethane solution was dried (Na₂SO₄), filtered and evaporated in vacuo.

A solution of thiol **8** (110 mg, 0.61 mmol) in dichloromethane (3 mL) was added to the crude epoxide (**9**) by cannula, rinsing with 1.2 mL. Basic alumina (628 mg) was added and the mixture stirred for 16 h. The solvent was evaporated and purification of the residue by flash chromatography on silica gel (15% ethyl acetate/hexanes) gave 129 mg (0.31 mmol, 48 %) of the coupled enone (**10**).

(3-Chloro-benzo[b]thiophen-2-yl)-methanol (12). To an ice cold solution of 10.0 g (47.0 mmol) of 3-chloro-benzo[b]thiophene-2-carboxylic acid (**11**) in 200 mL of THF was added 47 mL of LiAlH₄ (47 mmol, 1 M/THF). After 3 h, the reaction was quenched by addition of MeOH (ca. 40 mL). The volatiles were evaporated and the residue was treated with 50 mL 1 M HCl. After stirring for 10 min., the mixture was extracted with CH₂Cl₂ (3 x 150 mL). The combined CH₂Cl₂ solution was dried (MgSO₄), filtered and evaporated. Purification by flash chromatography on silica gel (10-20% ethyl acetate/hexane) gave 4.32 g (21.6 mmol, 46 %) of the alcohol (**12**).

3-Chloro-benzo[b]thiophene-2-carbaldehyde (13). A solution of alcohol **12** (4.32 g, 21.6 mmol) in 40 mL of CH₂Cl₂ was treated with 4A molecular sieves, NMO (3.81 g, 32.5 mmol), and TPAP (381 mg, 1.08 mmol). The reaction was stirred for 10 min. and then was evaporated to dryness. Purification by flash chromatography on silica gel (2% ethyl acetate/hexane) gave 3.52 g (18.3 mmol, 84%) of the aldehyde (**13**).

(E)-3-(3-Chloro-benzo[b]thiophen-2-yl)-acrylic acid methyl ester (14). A solution of 3.52 g (18.3 mmol) of **13** in 50 mL toluene was treated with methyl(triphenylphosphoranylidene)acetate (7.48 g, 21.9 mmol). After 4 h,

saturated NaHCO_3 solution (50 mL) was added and the mixture extracted with ethyl acetate (2 x 75 mL). The combined ethyl acetate solution was washed with brine (50 mL), dried (Na_2SO_4), filtered and evaporated. Purification by flash chromatography on silica gel (5% ethyl acetate/hexane) provided 3.60 g (14.6 mmol, 80%) of the enoate (**14**).

3-(3-Chloro-benzo[b]thiophen-2-yl)-propionic acid methyl ester (15). A solution of 3.60 g (14.6 mmol) of **14** in 50 mL THF was treated with Wilkinson's catalyst (3.35 g, 3.62 mmol). The mixture was stirred under 1 atm H_2 for 18 h and then was filtered through celite. The solvent was evaporated and the residue purified by flash chromatography on silica gel (0-2% ethyl acetate/hexane) to give 3.63 g (14.3 mmol, 99%) of the saturated ester (**15**).

3-(3-Chloro-benzo[b]thiophen-2-yl)-propan-1-ol (16). An ice cold solution of 3.63 g (14.3 mmol) of **15** in 60 mL of ether was treated with LiBH_4 (621 mg, 28.5 mmol) and methanol (2 mL). After 30 min., 30 mL of 0.5 M NaOH solution was added. The mixture was extracted with ethyl acetate (2 x 25 mL) and the combined ethyl acetate solution washed with brine (50 mL), dried (MgSO_4), filtered and evaporated. The residue was purified by flash chromatography on silica gel (5-20% ethyl acetate/hexane) to give 2.57 g (11.3 mmol, 79%) of the alcohol (**16**).

3-(3-Chloro-benzo[b]thiophen-2-yl)-propionaldehyde (17). A -78°C solution of oxalyl chloride (1.73 g, 13.6 mmol) in dichloromethane (20 mL) was treated with DMSO (20 mL). After 5 min., a solution of alcohol **16** (2.57g, 11.3 mmol) in dichloromethane (20 mL) was added. After another 15 min., triethylamine (7.1 mL, 50.6 mmol) was added. The reaction was stirred at -78°C

°C for 5 min., and then allowed to warm to room temperature. After 30 min., 100 mL water was added and the mixture extracted with dichloromethane (3 x 60 mL). The combined dichloromethane solution was dried (Na₂SO₄), filtered and evaporated. Purification by flash chromatography on silica gel (10% ethyl acetate/hexane) gave 2.11 g (9.4 mmol, 83%) of the aldehyde (**17**).

5-(3-Chloro-benzo[b]thiophen-2-yl)-pent-1-yn-3-ol (18). A solution of aldehyde **17** (2.11 g, 9.4 mmol) in 15 mL THF was added to a solution of ethynylmagnesium bromide (28.2 mL, 14.1 mmol, 0.5 M THF) at 0 °C. After 1.5 h, saturated NH₄Cl solution (75mL) was added and the mixture was extracted with ethyl acetate (3 x 50 mL). The combined ethyl acetate solution was washed with brine (50 mL) and then was dried (Na₂SO₄), filtered and evaporated. Purification by flash chromatography (5-20% ethyl acetate/hexane) gave 2.20 g (8.78 mmol, 93%) of the alcohol (**18**).

tert-Butyl-{1-[2-(3-chloro-benzo[b]thiophen-2-yl)-ethyl]-prop-2-ynyloxy}-dimethyl-silane (19). A solution of alcohol **18** (2.20 g, 8.78 mmol) in dichloromethane (15 mL) was treated with DMAP (215 mg, 1.8 mmol), TBSCl (1.59 g, 10.5 mmol), and triethylamine (1.8 mL, 13.2 mmol). The reaction was stirred for 24 h and then saturated sodium bicarbonate solution (50 mL) was added. The mixture was extracted with dichloromethane (2 x 50 mL) and the combined dichloromethane solution dried (Na₂SO₄), filtered and evaporated. Purification by flash chromatography (4% ethyl acetate/hexane) gave 3.06 g (6.4 mmol, 73%) of the protected alcohol (**19**).

(3-{{(1R,4S,5S)-4-(tert-Butyl-dimethyl-silanyloxy)-5-[(E)-3-(tert-butyl-dimethyl-silanyloxy)-5-(3-chloro-benzo[b]thiophen-2-yl)-pent-1-enyl]-3,3-

dimethyl-2-oxo-cyclopentylsulfanyl}-propylsulfanyl)-acetic acid methyl ester (20). A solution of alkyne **19** (105 mg, 0.28 mmol) in THF (1.2 mL) was treated with bis(cyclopentadienyl)zirconium chloride hydride (91 mg, 0.35 mmol). The reaction was stirred for 30 min., then was cooled to -78°C and
5 treated with methyllithium (0.46 mL, 0.64 mmol, 1.4 M in ether). After 10 min., a precooled (-78°C) solution of lithium 2-thienylcyanocuprate (1.3 mL, 0.33 mmol, 0.25 M in THF) was added by cannula. The reaction was stirred for 45 min. and then enone **10** (61 mg, 0.15 mmol) in 0.2 mL THF was added by cannula, rinsing with 0.2 mL THF. After 1 h, The reaction was quenched by
10 addition of 20 mL 1:1 saturated ammonium chloride solution/concentrated ammonium hydroxide. The mixture was stirred for 45 min. and then was extracted with ethyl acetate (3 x 20 mL). The combined ethyl acetate solution was dried (Na_2SO_4), filtered and evaporated. Purification by flash chromatography on silica gel (10 % ethyl acetate/hexanes) gave 51 mg (0.064
15 mmol, 43%) of the coupled product (**20**).

(3-((1R,4S,5S)-5-(3-chloro-benzo[b]thiophen-2-yl)-3-hydroxy-pent-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl}-propylsulfanyl)-acetic acid methyl ester (21, 22). A solution of **20** (51 mg, 0.064 mmol) in CH_3CN (1.6 mL) was treated with HF-pyridine (0.26 mL). The reaction was stirred for
20 24 h and then was quenched by addition of 15 mL saturated sodium bicarbonate solution. The mixture was extracted with dichloromethane (3 x 10 mL) and the combined dichloromethane solution was dried (Na_2SO_4), filtered and evaporated. Purification by preparative thin layer chromatography on silica gel

(40% ethyl acetate/hexanes) gave 12 mg (0.023 mmol, 71%) of each diastereomer.

(3-((1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid (23, 24). Rabbit liver esterase (9 mg) was added to a solution of the lower R_f ester **21** (11 mg, 0.021 mmol) in pH 7.2 phosphate buffer (0.5 mL)/CH₃CN (0.1 mL). The mixture was stirred overnight and then 10 mL 0.5 M HCl was added along with a few mL's of brine. The mixture was extracted with ethyl acetate (3 x 10 mL) and the combined ethyl acetate solution dried (Na₂SO₄), filtered and evaporated. Purification by flash chromatography on silica gel (3-5% MeOH/CH₂Cl₂) gave 4 mg (0.0078 mmol, 37%) of the acid (**23**). 300 MHz ¹H NMR (CDCl₃, ppm) δ 7.73 (2 H, d, J = 8.4 Hz) 7.4-7.3 (2 H, m) 5.9-5.8 (1 H, m) 5.8-5.7 (1 H, m) 4.4-4.3 (1 H, m) 3.63 (1 H, d, J = 9.7 Hz) 3.21 (2 H, s) 3.1-2.4 (11 H, overlapping m) 2.1-1.7 (4 H, overlapping m) 1.12 (3 H, s) 1.03 (3 H, s).

The higher R_f ester was hydrolyzed similarly except a solution of rabbit liver esterase (10 mg) in 0.5 mL of pH 7.2 phosphate buffer was added to a solution of the ester (10 mg, 0.019 mmol) in CH₃CN (0.2 mL). The reaction was stirred for 22 h and then worked up and purified as above. This gave 7 mg (0.013 mmol, 71%) of the acid (**24**). 300 MHz ¹H NMR (CDCl₃, ppm) δ 7.73 (2 H, d, J = 8.8 Hz) 7.44-7.31 (2 H, m) 5.9-5.8 (1 H, m) 5.8-5.7 (1 H, m) 4.4-4.3 (1 H, m) 3.64 (1 H, d, J = 9.7 Hz) 3.3-2.3 (13 H, overlapping m) 2.1-1.7 (4 H, overlapping m) 1.12 (3 H, s) 1.03 (3 H, s).

***tert*-Butyl-hex-5-ynyloxy-dimethyl-silane (26).**

7-(*tert*-Butyl-dimethyl-silanyloxy)-hept-2-yn-1-ol (27).**Acetic acid 7-(*tert*-butyl-dimethyl-silanyloxy)-hept-2-ynyl ester (28).** A

solution of 7-(*tert*-Butyl-dimethyl-silanyloxy)-hept-2-yn-1-ol **27** (4.507 g, 21 mmol) in pyridine (20 mL) was treated with acetic anhydride (3.0 mL, 31.8 mmol). After 18 h, the solvent was evaporated and the residue co-evaporated with toluene. The residue was used directly in the next step.

7-Acetoxy-hept-5-ynoic acid (29). A solution of crude **28** in acetone (100 mL)

was treated with Jones Reagent (18.0 mL, 41.4 mmol, 2.3 M). The mixture became warm and so was cooled with an ice bath. After 1 h at room

temperature, 10 mL isopropyl alcohol was added and the mixture stirred further for 15 min. The mixture still had a brown color so another 10 mL isopropyl alcohol was added. After another 15 min., the color had not changed so the mixture was filtered through celite and the filtrate evaporated in vacuo. The residue was partitioned between 100 mL ether and 100 mL saturated ammonium chloride solution. The aqueous layer was extracted with 100 mL ether and the combined ether solution washed with brine and then was dried (MgSO₄), filtered and evaporated to leave a yellow oil (6.333 g) that was used directly in the next step.

7-Hydroxy-hept-5-ynoic acid methyl ester (30). The crude acid **29** (6.333 g)

was treated with a 1% solution of acetyl chloride in methanol (60 mL). After 16 h, sodium bicarbonate (1.966 g, 23.4 mmol) was added. The mixture was dried (MgSO₄), filtered through celite and evaporated in vacuo. Purification by flash chromatography on silica gel (30-40% ethyl acetate/hexanes) gave 7-Hydroxy-

hept-5-ynoic acid methyl ester **30** (3.022 g, 19.3 mmol, 92% from 7-(*tert*-Butyl-dimethyl-silanyloxy)-hept-2-yn-1-ol **27**).

7-Iodo-hept-5-ynoic acid methyl ester (31). A solution of **30** (1.347 g, 8.6 mmol) in 5 mL dichloromethane was added to a mixture of triphenylphosphine
5 (2.725 g, 10.4 mmol), imidazole (726 mg, 10.7 mmol), and iodine (2.602 g, 10.3 mmol) in 34 mL dichloromethane, rinsing with 5 mL dichloromethane. After 40 min., the dichloromethane was evaporated in vacuo to a few mL's and the resulting mixture filtered through basic alumina, washing with 10% ethyl acetate/hexanes. Purification by flash chromatography on silica gel (10% ethyl
10 acetate/hexanes) gave 1.878 g (7.1 mmol, 83%) of the propargyl iodide.

***tert*-Butyl-[(*E*)-1-[2-(3-chloro-benzo[*b*]thiophen-2-yl)-ethyl]-3-iodo-allyloxy]-dimethyl-silane (32).** A solution of alkyne **19** (5.547 g, 15.2 mmol) in dichloromethane (50 mL) was treated with Cp₂ZrHCl (5.794 g, 22.5 mmol). The reaction was stirred for 45 min. and then N-iodosuccinimide (4.966 g, 22.1
15 mmol) was added. After 15 min., saturated sodium bicarbonate solution (200 mL) was added and the mixture was extracted with dichloromethane (2 x 100 mL). The combined dichloromethane solution was dried (MgSO₄), filtered and evaporated. Purification by flash chromatography on silica gel (0-5% ethyl acetate/hexanes) gave 6.608 g (13.1 mmol, 86%) of the vinyl iodide (**32**).

20 **7-[(1*R*,4*S*,5*R*)-4-(*tert*-Butyl-dimethyl-silanyloxy)-5-[(*E*)-3-(*tert*-butyl-dimethyl-silanyloxy)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-pent-1-enyl]-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (33).** A -78 °C solution of iodide **32** (675 mg, 1.34 mmol) in THF (2.0 mL) was treated with *tert*-butyllithium (1.73 mL, 2.94 mL, 1.7 M/pentane). The dark red mixture was

stirred for 25 min. and then dimethylzinc (0.80 mL, 1.6 mmol, 2 M/toluene) was added. The solution was stirred at 0 °C for 15 min. and then recooled to -78 °C. At this time, a solution of enone **6** (208 mg, 0.87 mmol) in THF (1.0 mL) was added over 2 h by syringe pump, rinsing with 0.5 mL THF. After 30 min.,

5 HMPA (1.34 mL, distilled from CaH₂) was added followed by a solution of propargyl iodide **31** (1.286 g, 4.83 mmol) in THF (1.0 mL). The solution was stirred in a -40 °C bath overnight and then 20 mL saturated ammonium chloride solution and 10 mL water were added. The mixture was extracted with dichloromethane (20 mL) and ethyl acetate (2 x 20 mL). The combined organic

10 extracts were dried (MgSO₄), filtered and evaporated. Purification by flash chromatography on silica gel (5-10% ethyl acetate/hexanes) gave 198 mg (0.27 mmol, 31%) of **33**.

(Z)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl}-hept-5-ynoic acid

15 **methyl ester (34, 35)**. A solution of **33** (198 mg, 0.27 mmol) in CH₃CN (6.5 mL) was treated with HF-pyridine (1.2 mL). The solution was stirred for 3 h and saturated sodium bicarbonate solution (120 mL) was added. The mixture was extracted with dichloromethane (3 x 50 mL) and the combined dichloromethane solution dried (Na₂SO₄), filtered and evaporated. Purification by flash

20 chromatography (50% ethyl acetate/hexane) followed by preparative TLC (55% ethyl acetate/hexane) gave 55 mg (0.11 mmol, 41%) of the less polar diastereomer (**34**) and 51 mg (0.10 mmol, 37%) of the more polar diastereomer (**35**).

(*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (low *R_f* diastereomer, **36**). A solution of **34** (9 mg, 0.017 mmol) and rabbit liver esterase (1 mg) in pH 7.2 phosphate buffer (2 mL)/CH₃CN (0.1 mL) was stirred for 17 h. The mixture was then coevaporated with CH₃CN to remove water and the residue purified by flash chromatography on silica gel (3-7% MeOH/CH₂Cl₂) to give 8 mg (0.016 mmol, 93%) of the acid (**36**).

(*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (high *R_f* diastereomer, **37**). A solution of **35** (12 mg, 0.023 mmol) and rabbit liver esterase (1 mg) in pH 7.2 phosphate buffer (2 mL)/CH₃CN (0.1 mL) was stirred for 17 h. TLC showed the presence of starting material, so another 2 mg of the esterase was added. After stirring for another 24 h, the reaction was complete. Work up and purification as above for **36** gave 8 mg (0.016 mmol, 69%) of the acid (**37**).

(*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (low *R_f* diastereomer, **38**). Ethanol (95%, 2.5 mL) was added to NiCl₂ (50 mg, 0.39 mmol) and NaBH₄ (7 mg, 0.19 mmol). The resulting black mixture was stirred for 5 min. and then ethylenediamine (41 µL, 0.61 mmol) was added. After 15 min., a solution of alkyne **34** (40 mg, 0.077 mmol) in 0.5 mL 95% ethanol was added, rinsing with 0.5 mL ethanol. The flask was purged with H₂ and allowed to stir under 1 atm H₂ for 22 h. The mixture was then

filtered through celite and purified by flash chromatography on silica gel (55% ethyl acetate/hexanes) to give 17 mg (0.032 mmol, 43%) of the alkene (**38**).

(Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid

5 **methyl ester (high R_f diastereomer 39)**. The same procedure as for **36** was followed to give 17 mg (0.032 mmol, 41%) of **39**.

(Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid

(low R_f diastereomer, **40**). The same procedure as above for **36** was used to
10 give 9 mg (0.018 mmol, 85%) of acid **40**. 300 MHz ¹H NMR (CDCl₃, ppm) δ
7.73 (2 H, d, J = 8.4 Hz) 7.45-7.30 (2 H, m) 5.8-5.6 (2 H, m) 5.4-5.3 (2 H, m)
4.3-4.1 (1 H, m) 3.57 (1 H, d, J = 9.7 Hz) 3.1-2.9 (2 H, m) 2.5-1.9 (10 H, m)
1.7-1.6 (2 H, m) 1.09 (3 H, s) 0.89 (3 H, s).

(Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid

15 **(high R_f diastereomer, 41)**. The same procedure as above for the **36** was used to give 9mg (0.018 mmol, 85%) of acid **41**. 300 MHz ¹H NMR (CDCl₃, ppm) δ
7.73 (2 H, d, J = 8.8 Hz) 7.45-7.30 (2 H, m) 5.8-5.6 (2 H, m) 5.45-5.30 (2 H, m)
4.3-4.2 (1 H, m) 3.61 (1 H, d, J = 9.7 Hz) 3.1-3.0 (2 H, m) 2.5-1.9 (10 H, m)
20 1.7-1.6 (2 H, m) 1.10 (3 H, s) 0.90 (3 H, s).

(Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (2-hydroxy-

ethyl)-amide (65). A solution of acid **55** (prepared similarly to acid **41**, 7 mg, 0.015 mmol) in DMF (0.5 mL) was treated with N-hydroxysuccinimide (6.9 mg,

0.056 mmol). The mixture was stirred for 5 minutes and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDCI, 20.7 mg, 0.11 mmol) was added. After stirring for 7 h, 2-aminoethanol (5 μ L, 0.083 mmol) was added and the mixture stirred further for 16 h. Ethyl acetate (50 mL) was added and the mixture was washed with water (3 x 50 mL) and brine (50 mL). The organic layer was dried (Na_2SO_4), filtered and evaporated. Purification by flash chromatography on silica gel (5% methanol/dichloromethane) followed by preparative thin layer chromatography (10% methanol/dichloromethane) gave amide **65** (5 mg, 0.010 mmol, 65%).

10 **(Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid amide (69).** A solution of acid **55** (9 mg, 0.02 mmol) in dichloromethane (0.2 mL) was treated with triethylamine (15 μ L, 0.11 mmol). The solution was cooled to 0 °C and after 10 minutes, ethyl chloroformate (7 μ L, 0.073 mmol) was added. The

15 solution was stirred further for 1 h at 0 °C and then concentrated aqueous ammonium hydroxide solution was added (10 μ L, 0.26 mmol). The reaction was allowed to stir at room temperature overnight and then was quenched by addition of 0.5 M HCl (7 mL). The mixture was extracted with ethyl acetate (3 x 30 mL) and the combined ethyl acetate solution was washed with saturated

20 NaHCO_3 solution (20 mL) and brine (20 mL) and then was dried (Na_2SO_4), filtered and evaporated. Purification by flash chromatography on silica gel (2% - 6% methanol/dichloromethane) gave the title amide (2.6 mg, 28 %).

The methods of screening the compounds of this invention for the desired biological activity are illustrated in the following non-limiting examples. Results for example compounds of this invention are included in Table 2. These results are presented purely for illustrative purposes and are not intended to limit the scope of the invention in any way.

Radioligand Binding

Cells Stably Expressing EP₁, EP₂, EP₄ and FP Receptors

HEK-293 cells stably expressing the human or feline FP receptor, or EP₁, EP₂, or EP₄ receptors were washed with TME buffer, scraped from the bottom of the flasks, and homogenized for 30 sec using a Brinkman PT 10/35 polytron. TME buffer was added to achieve a final 40 ml volume in the centrifuge tubes (the composition of TME is 100 mM TRIS base, 20 mM MgCl₂, 2M EDTA; 10N HCl is added to achieve a pH of 7.4).

The cell homogenate was centrifuged at 19000 r.p.m. for 20 min at 4° C using a Beckman Ti-60 rotor. The resultant pellet was resuspended in TME buffer to give a final 1 mg/ml protein concentration, as determined by Biorad assay. Radioligand binding competition assays vs. [³H]-17-phenyl PGF_{2α} (5 nM) were performed in a 100μl volume for 60 min. Binding reactions were started by adding plasma membrane fraction. The reaction was terminated by the addition of 4 ml ice-cold TRIS-HCl buffer and rapid filtration through glass fiber GF/B filters using a Brandel cell harvester. The filters were washed 3 times with ice-cold buffer and oven dried for one hour.

[³H]-PGE₂ (specific activity 180 Ci mmol) was used as the radioligand for EP receptors. [³H] 17-phenyl PGF_{2α} was employed for FP receptor binding studies. Binding studies employing EP₁, EP₂, EP₄ and FP receptors were performed in duplicate in at least three separate experiments. A 200μl assay volume was used. Incubations were for 60 min at 25°C and were terminated by the addition of 4 ml of ice-cold 50 mM TRIS-HCl, followed by rapid filtration through Whatman GF/B filters and three additional 4 ml washes in a cell harvester (Brandel). Competition studies were performed using a final concentration of 5 nM [³H]-PGE₂, or 5 nM [³H] 17-phenyl PGF_{2α} and non-

specific binding determined with 10^{-5} M of unlabeled PGE₂, or 17-phenyl PGF_{2 α} , according to receptor subtype studied.

METHODS FOR FLIPR™ STUDIES

(a) CELL CULTURE

5 HEK-293(EBNA) cells, stably expressing one type or subtype of recombinant human prostaglandin receptors (prostaglandin receptors expressed: hDP/Gqs5; hEP₁; hEP₂/Gqs5; hEP_{3A}/Gqi5; hEP₄/Gqs5; hFP; hIP; hTP), were cultured in 100 mm culture dishes in high-glucose DMEM medium containing 10% fetal bovine serum, 2 mM l-glutamine, 250 µg/ml geneticin (G418) and
10 200 µg/ml hygromycin B as selection markers, and 100 units/ml penicillin G, 100 µg/ml streptomycin and 0.25 µg/ml amphotericin B.

(b) CALCIUM SIGNAL STUDIES ON THE FLIPR™

Cells were seeded at a density of 5×10^4 cells per well in Biocoat® Poly-D-lysine-coated black-wall, clear-bottom 96-well plates (Becton-Dickinson) and
15 allowed to attach overnight in an incubator at 37 °C. Cells were then washed two times with HBSS-HEPES buffer (Hanks Balanced Salt Solution without bicarbonate and phenol red, 20 mM HEPES, pH 7.4) using a Denley Cellwash plate washer (Labsystems). After 45 minutes of dye-loading in the dark, using the calcium-sensitive dye Fluo-4 AM at a final concentration of 2 µM, plates
20 were washed four times with HBSS-HEPES buffer to remove excess dye leaving 100 µl in each well. Plates were re-equilibrated to 37 °C for a few minutes.

Cells were excited with an Argon laser at 488 nm, and emission was measured through a 510-570 nm bandwidth emission filter (FLIPR™,
25 Molecular Devices, Sunnyvale, CA). Drug solution was added in a 50 µl volume to each well to give the desired final concentration. The peak increase in fluorescence intensity was recorded for each well. On each plate, four wells each served as negative (HBSS-HEPES buffer) and positive controls (standard agonists: BW245C (hDP); PGE₂ (hEP₁; hEP₂/Gqs5; hEP_{3A}/Gqi5; hEP₄/Gqs5);
30 PGF_{2 α} (hFP); carbacyclin (hIP); U-46619 (hTP), depending on receptor). The

peak fluorescence change in each drug-containing well was then expressed relative to the controls.

Compounds were tested in a high-throughput (HTS) or concentration-response (CoRe) format. In the HTS format, forty-four compounds per plate were examined in duplicates at a concentration of 10^{-5} M. To generate concentration-response curves, four compounds per plate were tested in duplicates in a concentration range between 10^{-5} and 10^{-11} M. The duplicate values were averaged. In either, HTS or CoRe format each compound was tested on at least 3 separate plates using cells from different passages to give an $n \geq 3$.

10

Table 2

Compound	hFP	hEP ₁	hEP ₂	hEP _{3D} hEP _{3A}	hEP ₄	hDP	hIP	hTP
21	NA	NA	>10 K	NA	98	NA	NA	NA
22	NA	NA	NA NA	NA NA	300 30	NA	NA	NA
23	NA	NA	NA NA	>10K NA	44 0.1	NA	NA	>10K
24	NA	NA	NA NA	>>10K NA	26 0.1	NA	NA	NA
34	NA		NA	NA	>10K >10K		NA	NA
35	NA			NA	2455		NA	NA
36	NA		NA	NA	200 66		>10K	NA
37	NA		NA	NA	100 32		>10K	NA
38	NA		NA	NA	2700 269		NA	NA
39	NA		NA	NA	2300 141		NA	NA
40	NA		NA	NA	200 0.3		NA	>10K
41	NA		>10K	NA	20		NA	>10K
42	NA	NA	NA NA	>10 ⁴ 559	>10 ⁴ NA	NA	NA	NA
43	NA	>10 ⁴	NA NA	1700 11	400 63		3981	18
44	NA	782	1500 944	300 4.6	5.5 0.2	>10 K	284	18
45	NA	NA	NA NA	>10 ⁴ 531	400 51	NA	NA NA	631 NA
46	NA	290	>10K	>10K 589	4 0.4	NA	NA	

47	NA	963	NA	>10K	76		NA	
48			NA		45			
49			NA		1400			
50	NA	638	NA	6607 >10K	2400 3162		NA	>10K
51	NA		NA	NA	700		NA	
52	NA	27	NA	60	72 18		NA	
53	NA	1020	NA	1862	59 6.4		NA	
54	NA	308	NA NA	4700	20 0.3	NA	NA	
55	NA	758	NA NA	>10K	310 38	NA	NA	
60	NA	NA	NA	NA	>10K	NA	NA	NA
61	NA	NA	NA	NA	NA	NA	NA	NA
62	NA	NA	NA	NA	832	NA	NA	NA
63	NA	>10K	NA	NA	478	NA	NA	NA
64	NA	NA	NA	NA	4154	NA	NA	NA
65	NA	NA	NA	NA	NA	NA	NA	NA
68	NA	NA	NA	NA	678	NA	NA	>10K
69	NA	NA	NA	NA	5000	NA	NA	>10K
70	NA	NA	NA	>10K	219	NA	NA	
71	NA	NA	NA	NA	10000	NA	NA	
72	NA	NA	NA	NA	>10K	NA	NA	NA
73	NA	NA	NA	NA	NA	NA	NA	NA
74	NA	2376	NA		256	NA	NA	
75	NA	2050	NA		>10K	NA	NA	>10K

The top numbers are the radioligand binding values(nm)

The bottom numbers are the functional data (nm)

5 The foregoing description details specific methods and compositions that can be employed to practice the present invention, and represents the best mode contemplated. However, it is apparent for one of ordinary skill in the art that


further compounds with the desired pharmacological properties can be prepared in an analogous manner, and that the disclosed compounds can also be obtained from different starting compounds via different chemical reactions. Similarly, different pharmaceutical compositions may be prepared and used with

5 substantially the same result. Thus, however detailed the foregoing may appear in text, it should not be construed as limiting the overall scope hereof; rather, the ambit of the present invention is to be governed only by the lawful construction of the appended claims.

CLAIMS

What is claimed is:

1. Use of a compound of Formula I for the manufacture of a medicament
5 for treating ocular hypertension or glaucoma in a mammal,



Formula I

10

wherein the dashed lines indicate the presence or absence of a bond, the hatched wedges indicate the α (down) configuration, and the solid triangles indicate the β (up) configuration;

B is a single, double, or triple covalent bond;

15


n is 0-6;

X is CH₂, S or O;

Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂,

CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, CH₂OR, P(O)(OR)₂, CONRSO₂R,

SONR₂, or



20

R is H, C₁₋₆ alkyl or C₂₋₆ alkenyl;


R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached;

R^4 is hydrogen, R, $C(=O)R$, or any group that is easily removed under physiological conditions such that R^4 is effectively hydrogen;

R^5 is hydrogen or R; and

R^6 is

- 5 i) hydrogen;
 - ii) a linear or branched hydrocarbon containing between 1 and 8 carbon atoms, which may contain one or more double or triple bonds, or oxygen or halogen derivatives of said hydrocarbon, wherein 1-3 carbon or hydrogen atoms may be substituted by O or a halogen; or
 - 10 iii) aryloxy, heteroaryloxy, C_{3-8} cycloalkyloxy, C_{3-8} cycloalkyl, C_{6-10} aryl or C_{3-10} heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{6-10} aryl, C_{3-10} heteroaryl, aryloxy, heteroaryloxy, C_{1-6} alkyl,
 - 15 OR, SR, and SO_2R .
2. Use according to claim 1 wherein the compound of Formula I is not a compound of Formula II




Formula II

- 20 wherein A is CO_2H or CO_2Me ;
- D is a single, double, or triple covalent bond;
- E is a linear, branched, or cycloalkyl chain of 3 to 7 carbons, trifluoromethylbutyl, hydroxylalkyl, or CH_2R^7 wherein R^7 is phenyl, cyclopentyl, phenoxy, chlorophenoxy, propoxy, or $-CH_2SCH_2CH_3$;

J is hydrogen, R, C(=O)R, or any group that is easily removed under physiological conditions such that R⁴ is effectively hydrogen; and G is H or CH₃.


3. Use according to claim 2 wherein the compound of Formula I is further represented by Formula III



Formula III

wherein Y is CO₂R, or any pharmaceutically acceptable salt of CO₂H.

4. Use according to claim 3 wherein R⁶ is C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.
5. Use according to claim 4 wherein R⁶ is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.
6. Use according to claim 5 wherein Y is CO₂H or CO₂Me.
7. Use according to claim 6 wherein R⁶ is 3-chlorobenzothien-2-yl.
8. Use according to claim 7 wherein n is 2.
9. Use according to claim 8 wherein B is a single bond.
10. Use according to claim 2 wherein the compound of Formula I is further represented by Formula IV

**Formula IV**

wherein Y is CO₂R, or any pharmaceutically acceptable salt of CO₂H; and

R⁶ is C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.

11. Use according to claim 10 wherein Y is CO₂H or CO₂Me.

12. Use according to claim 11 wherein R⁶ is phenyl.

13. Use according to claim 12 wherein B is a double bond.

10 14. Use according to claim 11 wherein R⁶ is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.

15. Use according to claim 14 wherein R⁶ is 3-chlorobenzothien-2-yl.

15 16. Use according to claim 15 wherein B is a double or triple bond.


17. Use according to claim 1 wherein said compound is selected from the group consisting of

(3-((1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid methyl ester (**21**, **22**);

20 (3-((1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid (**23**, **24**);

- (Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**34, 35**);
- (Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**36,37**);
- 5 (Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**38,39**);
- (Z)-7-[(1R,4S,5R)-5-[(E)-5-(3-chloro-benzo[b]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**40,41**);
- 10 7-[(1R,4S,5R)-4-Hydroxy-5-((E)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**42**)
- 7-[(1R,4S,5R)-4-Hydroxy-5-((E)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**43**)
- 15 (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**44**)
- (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-(S)-3-hydroxy-oct-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**45**)
- (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-4-phenyl-but-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**46, 47**)
- 20 (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-4-phenyl-but-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**48, 49**)
- (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**50,51**)
- 25 (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**52,53**)
- (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**54,55**)
- 7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-heptanoic acid (**56,57**)
- 30 (Z)-7-[(1R,4S,5R)-5-(4-Benzo[b]thiophen-2-yl-3-hydroxy-butyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**58,59**)

- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid ethylamide (**60,61**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid diethylamide
- 5 (**62,63**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (2-hydroxy-ethyl)-amide (**64,65**)
- (3*S*,4*R*,5*R*)-4-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-3-hydroxy-
- 10 2,2-dimethyl-5-[(*Z*)-6-(1-*H*-tetrazol-5-yl)-hex-2-enyl]-cyclopentanone (**66,67**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid amide (**68,69**)
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**70,71**)
- 15 7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**72,73**)
- 7-[(1*R*,4*S*,5*R*)-5-((*E*)-4-Benzo[*b*]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**74,75**)
18. A compound represented by Formula I:



Formula I

20

wherein the dashed lines indicate the presence or absence of a bond, the hatched wedges indicate the α (down) configuration, and the solid triangles indicate the

25 β (up) configuration;


B is a single, double, or triple covalent bond;

n is 0-6;

X is CH₂, S or O wherein both X moieties are identical;

Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂,

- 5 CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, CH₂OR, P(O)(OR)₂, CONRSO₂R, SONR₂, or



R is H, C₁₋₆ alkyl or C₂₋₆ alkenyl;

- 10 R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached;

R⁴ is hydrogen, R, C(=O)R, or any group that is easily removed under physiological conditions such that R⁴ is effectively hydrogen;


R⁵ is hydrogen or R;

- 15 R⁶ is

- i) hydrogen;
- ii) a linear or branched hydrocarbon containing between 1 and 8 carbon atoms, which may contain one or more double or triple bonds, or oxygen or halogen derivatives of said hydrocarbon, wherein 1-3 carbon or hydrogen atoms may be substituted by O or a halogen; or
- 20 iii) aryloxy, heteroaryloxy, C₃₋₈ cycloalkyloxy, C₃₋₈ cycloalkyl, C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino,
- 25 hydroxy, C₆₋₁₀ aryl, C₃₋₁₀ heteroaryl, aryloxy, heteroaryloxy, C₁₋₆ alkyl, OR, SR, and SO₂R; and

the compound of Formula I is not a compound of Formula II

48

**Formula II**


wherein A is CO₂H, CO₂Me, or CO₂Et;

D is a single, double, or triple covalent bond;

- 5 E is a linear, branched, or cycloalkyl chain of 3 to 7 carbons, trifluoromethylbutyl, hydroxylalkyl, or CH₂R⁷ wherein R⁷ is phenyl, cyclopentyl, phenoxy, chlorophenoxy, propoxy, or -CH₂SCH₂CH₃;
- J is hydrogen, R, C(=O)R, or any group that is easily removed under physiological conditions such that R⁴ is effectively hydrogen; and

- 10 G is H or CH₃.


19. The compound of claim 18 which is further represented by Formula III

**Formula III**

wherein Y is CO₂R, or any pharmaceutically acceptable salt of CO₂H.

20. The compound of claim 19 wherein R⁶ is C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl,
- 15 wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.


21. The compound of claim 20 wherein R^6 is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R .
- 5 22. The compound of claim 21 wherein Y is CO_2H or CO_2Me .
23. The compound of claim 22 where R^6 is 3-chlorobenzothien-2-yl.
24. The compound of claim 23 where n is 2.
25. The compound of claim 24 where B is a single bond.
- 10 26. The compound of claim 18 which is further represented by Formula IV



Formula IV

- wherein Y is CO_2R or any pharmaceutically acceptable salt of CO_2H ; and R^6 is C_{6-10} aryl or C_{3-10} heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R .
- 15 27. The compound of claim 26 wherein Y is CO_2H or CO_2Me .
28. The compound of claim 27 wherein R^6 is phenyl.
29. The compound of claim 28 wherein B is a double bond.
- 20 30. The compound of claim 27 wherein R^6 is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C_{1-6} alkyl, OR, SR, and SO_2R .
31. The compound of claim 30 wherein R^6 is 3-chlorobenzothien-2-yl.
- 25 32. The compound of claim 31 wherein B is a double or triple bond.

33. The compound of claim 18 which is further represented by Formula V




Formula V

wherein at least one of R^2 and R^3 is not methyl.

34. The compound of claim 33 wherein R^2 and R^3 have a total number of
5 carbon atoms of 6 or less.
35. The compound of claim 34 wherein R^5 is hydrogen.
36. The compound of claim 18 wherein said compound is selected from the
group consisting of
- 10 (3-[(1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid methyl ester (**21**, **22**);
- (3-[(1*R*,4*S*,5*S*)-5-(3-chloro-benzo[*b*]thiophen-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentylsulfanyl)-propylsulfanyl)-acetic acid
(**23**, **24**);
- 15 (*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**34**, **35**);
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**36**, **37**);
- 20 (*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**38**, **39**);
- (*Z*)-7-[(1*R*,4*S*,5*R*)-5-[(*E*)-5-(3-chloro-benzo[*b*]thiophene-2-yl)-3-hydroxy-pent-1-enyl]-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**40**, **41**);

- (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**50,51**)
- (Z)-7-[(1R,4S,5R)-4-Hydroxy-5-((E)-3-hydroxy-5-phenyl-pent-1-enyl)-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**52,53**)
- 5 (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**54,55**)
- 7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-heptanoic acid (**56,57**)
- (Z)-7-[(1R,4S,5R)-5-(4-Benzo[b]thiophen-2-yl-3-hydroxy-butyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (**58,59**)
- 10 (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid ethylamide (**60,61**)
- (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid diethylamide
- 15 (**62,63**)
- (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid (2-hydroxy-ethyl)-amide (**64,65**)
- (3S,4R,5R)-4-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-3-hydroxy-
- 20 2,2-dimethyl-5-[(Z)-6-(1-H-tetrazol-5-yl)-hex-2-enyl]-cyclopentanone (**66,67**)
- (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid amide (**68,69**)
- (Z)-7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-enoic acid methyl ester (**70,71**)
- 25 7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid methyl ester (**72,73**)
- 7-[(1R,4S,5R)-5-((E)-4-Benzo[b]thiophen-2-yl-3-hydroxy-but-1-enyl)-4-hydroxy-3,3-dimethyl-2-oxo-cyclopentyl]-hept-5-ynoic acid (**74,75**)
37. An ophthalmic solution comprising a therapeutically effective amount of a
- 30 compound represented by the general Formula 1

52

**Formula I**


- 5 wherein the dashed lines indicate the presence or absence of a bond, the hatched wedges indicate the α (down) configuration, and the solid triangles indicate the β (up) configuration;

B is a single, double, or triple covalent bond;

n is 0-6;

- 10 X is CH₂, S or O;

Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂, CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, CH₂OR, P(O)(OR)₂, CONRSO₂R, SONR₂, or



- 15 R is H, C₁₋₆ alkyl or C₂₋₆ alkenyl;

R² and R³ are C₁₋₆ linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached;

R⁴ is hydrogen, R, C(=O)R, or any group that is easily removed under

- 20 physiological conditions such that R⁴ is effectively hydrogen;

R⁵ is hydrogen or R; and


R⁶ is

- i) hydrogen;

- ii) a linear or branched hydrocarbon containing between 1 and 8 carbon atoms, which may contain one or more double or triple bonds, or oxygen or halogen derivatives of said hydrocarbon, wherein 1-3 carbon or hydrogen atoms may be substituted by O or a halogen; or
- 5 iii) aryloxy, heteroaryloxy, C₃₋₈ cycloalkyloxy, C₃₋₈ cycloalkyl, C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₆₋₁₀ aryl, C₃₋₁₀ heteroaryl, aryloxy, heteroaryloxy, C₁₋₆ alkyl,
- 10 OR, SR, and SO₂R.



38. A pharmaceutical product, comprising a container adapted to dispense the contents of said container in metered form; and an ophthalmic solution according to claim 37 in said container.

- 15 39. Use according to making compounds of Formula V



Formula V

- wherein part of the synthesis comprises reducing a compound of Formula VI to a
- 20 compound of Formula VII in the presence of Baker's yeast

**Formula VI****Formula VII**

wherein the dashed lines indicate the presence or absence of a bond, the hatched wedges indicate the α (down) configuration, and the solid triangles indicate the β (up) configuration;

5 B is a single, double, or triple covalent bond;


n is 0-6;

X is CH_2 , S or O;

Y is any pharmaceutically acceptable salt of CO_2H , CO_2R , CONR_2 ,

$\text{CONHCH}_2\text{CH}_2\text{OH}$, $\text{CON}(\text{CH}_2\text{CH}_2\text{OH})_2$, CH_2OR , $\text{P}(\text{O})(\text{OR})_2$, CONRSO_2R ,

10 SONR_2 , or



R is H, C_{1-6} alkyl or C_{2-6} alkenyl;

R^2 and R^3 are C_{1-6} linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to

15 which they are commonly attached;

R^5 is hydrogen or R; and

R^6 is


i) hydrogen;

ii) a linear or branched hydrocarbon containing between 1 and 8

20 carbon atoms, which may contain one or more double or triple bonds, or oxygen or halogen derivatives of said hydrocarbon, wherein 1-3 carbon or hydrogen atoms may be substituted by O or a halogen; or


iii) aryloxy, heteroaryloxy, C₃₋₈ cycloalkyloxy, C₃₋₈ cycloalkyl, C₆₋₁₀ aryl or C₃₋₁₀ heteroaryl, wherein one or more carbons is substituted with N, O, or S; and which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₆₋₁₀ aryl, C₃₋₁₀ heteroaryl, aryloxy, heteroaryloxy, C₁₋₆ alkyl, OR, SR, and SO₂R.

40. The method of making compounds of Formula VIII comprising:




Formula VIII

i) reacting a compound of Formula IX with a compound of Formula X in the presence of a suitable base to form a compound of Formula XI;




Formula IX




Formula X

56

**Formula XI**

ii) coupling a compound of Formula XI with a compound of Formula XII; and

**Formula XII**

5 iii) removing the protecting groups and separating the diastereomers to obtain the desired products;

wherein the hatched wedges indicate the α (down) configuration, the solid triangles indicate the β (up) configuration, and the wavy lines indicate either the cis (Z) or trans (Z) conformation;

10 n is 0-6;

B is a single, double, or triple covalent bond;

J is a protecting group that can be easily removed to form the respective hydroxide group without affecting the rest of the molecule;

R is C_{1-6} alkyl or C_{2-6} alkenyl;


15 R^2 and R^3 are C_{1-6} linear alkyl which may be the same or different, and may be bonded to each other such that they form a ring incorporating the carbon to which they are commonly attached;

X is S or O wherein both X moieties are identical; and

M is a group that comprises one or more metal atoms.

20 41. Use according to claim 1 wherein the compound of Formula I which is further represented by Formula XIII

57


**Formula XIII**

wherein B represents a single or double bond;

and R⁶ is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.

42. Use according to claim 41 wherein R⁶ is benzothien-2-yl.

43. Use according to claim 42 wherein Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂, CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, or




44. Use according to claim 43 wherein the dashed line indicates the presence of a bond and B is a double bond.

45. Use according to claim 43 wherein the dashed line indicates the presence of a bond and B is a single bond.

46. Use according to claim 43 wherein the dashed line indicates the absence of a bond and B is a double bond.

47. The compound of claim 18 which is further represented by Formula XIII

58


**Formula XIII**

wherein B represents a single or double bond;

and R⁶ is naphthyl, benzofuranyl, or benzothienyl, which may contain one or more substituents selected from the group consisting of halogen, trihalomethyl, cyano, nitro, amino, hydroxy, C₁₋₆ alkyl, OR, SR, and SO₂R.

48. The compound of claim 47 wherein R⁶ is benzothien-2-yl.

49. The compound of claim 48 wherein Y is any pharmaceutically acceptable salt of CO₂H, or CO₂R, CONR₂, CONHCH₂CH₂OH, CON(CH₂CH₂OH)₂, or



50. The compound of claim 49 wherein the dashed line indicates the presence of a bond and B is a double bond.


51. The compound of claim 49 wherein the dashed line indicates the presence of a bond and B is a single bond.

52. The compound of claim 49 wherein the dashed line indicates the absence of a bond and B is a double bond.


53. The compound of claim 18 wherein A is CO₂R⁸, wherein R⁸ is any linear, branched, or cyclic alkyl group having from 3 to 6 carbons.

FIG. 1.
(SCHEME 1)

EQUATION 1



EQUATION 2



EQUATION 3





FIG. 2.
(SCHEME 2)



2/5

FIG. 3.
(SCHEME 3)










FIG. 5.
(SCHEME 5)



3/5

Fig. 4.
(SCHEME 4)



4/5
FIG. 6.
(SCHEME 6)





FIG. 8.
(SCHEME 8)



5/5

FIG. 7.
(SCHEME 7)

